# Author Index


AALBERSE, R.C. & SCHUURMAN, J. IgG4 breaking the rules, 9
ACUNA, C.L. *see* ZHANG, Y.
AJUEBOR, M.N. & SWAIN, M. G. Role of chemokines and chemokine receptors in the gastrointestinal tract, 137
AKAO, S. *see* YAMAKAMI, K.
AKASHI, H. *see* SHIMOJIMA, M.
AKPORIAYE, E. *see* ROBEY, I.F.
ALEXANDER, I.E. *see* LOGAN, G.J.
AMPHLETT, G.E. *see* FENHALLS, G.
ANANABA, G.A. *see* MOORE, T.
ANDERSON, G. *see* HARE, K.J.

BALDWIN, C.L. *see* SATHIYASEELAN, T.
BARDIN, P. *see* FENHALLS, G.
BAR-SHAVIT, R., MAOZ, M., YONGJUN, Y., GROYSMAN, M., DEKEL, I. & KATZAV, S. Signalling pathways induced by protease-activated receptors and integrins in T cells, 35
BEIGNON, A.-S., BRIAND, J.-P., MULLER, S. & PARTIDOS, C. D. Immunization onto bare skin with synthetic peptides: immunomodulation with a CpG-containing oligodeoxynucleotide and effective priming of influenza virus-specific $CD4^+$ T cells, 204
BELAY, T. *see* MOORE, T.
BERKE, G. *see* LI, J.-H.
BERUMEN, J. *see* GARCÍA-HERNÁNDEZ, M.L.
BEZUIDENHOUT, J. *see* FENHALLS, G.
BICE, D.E. *see* OUT, T.A.
BILENKI, L. *see* YANG, X.
BLACK, S.J. *see* SATHIYASEELAN, T.
BOLIER, J. *see* MOORE, T.
BOWERS, S. *see* MOORE, T.
BRADLEY, D.S. *see* ELLISON, C.A.
BRIAND, J.-P. *see* BEIGNON, A.-S.
BROWN, W.C. *see* FRASER, D.G.

CANTRELL, D.A. T-cell antigen receptor signal transduction, 369
CAULFIELD, J., FERNANDEZ, M., SNETKOV, V., LEE, T. & HAWRYLOWICZ, C. CXCR4 expression on monocytes is up-regulated by dexamethasone and is modulated by autologous $CD3^+$ T cells, 155
CHEN, L. *see* YANG, X.
CHEN, Y., MA, Y. & CHEN, Y. Roles of cytotoxic T-lymphocyte-associated antigen-4 in the inductive phase of oral tolerance, 171
CHEN, Y. *see* CHEN, Y.
CHIASSON, R.J. *see* HINKLEY, K.S.
CHILDS, R.A. *see* GALUSTIAN, C.
CHOI, J.W. *see* JEONG, J.-Y.
CROMPTON, T. *see* OUTRAM, S.V.
CUMBERBATCH, M., DEARMAN, R.J. & KIMBER, I. Influence of ageing on Langerhans cell migration in mice: identification of a putative deficiency of epidermal interleukin-1β, 466
CURNOCK, A.P., LOGAN, M.K. & WARD, S. G. Chemokine signalling: pivoting around multiple phosphoinositide 3-kinases, 125

DAVIDSON, J. *see* ROTONDO, D.
DEARMAN, R.J. *see* CUMBERBATCH, M.
DEKEL, I. *see* BAR-SHAVIT, R.
DELVIG, A.A. *see* ROBINSON, J.H.
DIAO, H., KOHANAWA, M., YIMIN, H., NAKAJIMA, Y., SATO, T., MINAGAWA, A. & NAKANE, A. Lipopolysaccharide triggers invasive streptococcal disease in mice through a tumour necrosis factor-α-dependent mechanism, 344
DOYLE, C. *see* FAIOLA, B.
DUNCAN, K. *see* FENHALLS, G.
DURHAM, S.R. *see* WACHHOLZ, P.R.

EARL, J.W. *see* LOGAN, G.J.
EKO, F.O. *see* MOORE, T.
ELLISON, C.A., BRADLEY, D.S., FISCHER, J.M.M., HAYGLASS, K.T. & GARTNER, J. G. Murine graft-versus-host disease induced using interferon-γ-deficient grafts features antibodies to double-stranded DNA, T helper 2-type cytokines and hypereosinophilia, 63

FAIOLA, B., DOYLE, C., GILBOA, E. & NAIR, S. Influence of CD4 T cells and the source of major histocompatibility complex class II-restricted peptides on cytotoxic T-cell priming by dendritic cells, 47
FAN, Y. *see* YANG, X.
FAYYAZI, A. *see* SORURI, A.
FEIZI, T. *see* GALUSTIAN, C.
FENHALLS, G., STEVENS, L., BEZUIDENHOUT, J., AMPHLETT, G.E., DUNCAN, K., BARDIN, P. & LUKEY, P. T. Distribution of IFN-γ, IL-4 and TNF-α protein and CD8 T cells producing IL-12p40 mRNA in human lung tuberculous granulomas, 325
FERNANDEZ, M. *see* CAULFIELD, J.
FISCHER, J.M.M. *see* ELLISON, C.A.
FRACKER, P. *see* LAAKKO, T.
FRASER, D.G., OAKS, J.L., BROWN, W.C. & MCGUIRE, T. C. Identification of broadly recognized, T helper 1 lymphocyte epitopes in an equine lentivirus, 295
FUKUI, T. *see* IMAI, Y.

GALUSTIAN, C., CHILDS, R.A., STOLL, M., ISHIDA, H., KISO, M. & FEIZI, T. Synergistic interactions of the two classes of ligand, sialyl-Lewis$^{a/x}$ fuco-oligosaccharides and short sulpho-motifs, with the P- and L-selectins: implications for therapeutic inhibitor designs, 350
GARCÍA-HERNÁNDEZ, M.L., HERNÁNDEZ-PANDO, R., GARIGLIO, P. & BERUMEN, J. Interleukin-10 promotes B16-melanoma growth by inhibition of macrophage functions and induction of tumour and vascular cell proliferation, 231
GARIGLIO, P. *see* GARCÍA-HERNÁNDEZ, M.L.
GARTNER, J.G. *see* ELLISON, C.A.
GILBOA, E. *see* FAIOLA, B.
GILMORE, X. *see* ZHANG, Y.
GODDARD, E.A. *see* HUSSEY, G.D.
GOTTSCHALK, S. *see* HUSSEY, G.D.
GRAHAM, C.M. *see* SMITH, C.A.
GROYSMAN, M. *see* BAR-SHAVIT, R.

HÄCKER, G., REDECKE, V. & HÄCKER, H. Activation of the immune system by bacterial CpG-DNA, 245
HÄCKER, H. *see* HÄCKER, G.
HAN, X. *see* YANG, X.
HARE, K.J., PONGRACZ, J., JENKINSON, E.J. & ANDERSON, G. Induction of thymocyte positive selection does not convey immediate resistance to negative selection, 163

HARRIMAN, G.R. *see* ZHANG, Y.
HARRIS, J., WERLING, D., KOSS, M., MONAGHAN, P., TAYLOR, G. & HOWARD, C. J. Expression of caveolin by bovine lymphocytes and antigen-presenting cells, 190
HARRIS, S.G. & PHIPPS, R. P. Prostaglandin $D_2$, its metabolite 15-d-$PGJ_2$, and peroxisome proliferator activated receptor-$\gamma$ agonists induce apoptosis in transformed, but not normal, human T lineage cells, 23
HASHIMOTO, K. *see* UCHIYAMA, H.
HASKARD, D.O. *see* KHAN, A.I.
HAWRYLOWICZ, C. *see* CAULFIELD, J.
HAY, A. *see* SMITH, C.
HAYGLASS, K. *see* ELLISON, C.
HERNÁNDEZ-PANDO, R. *see* GARCÍA-HERNÁNDEZ, M.L.
HINKLEY, K.S., CHIASSON, R., PRIOR, T. & RIGGS, J. E. Age-dependent increase of peritoneal B-1b B cells in SCID mice, 196
HOWARD, C.J. *see* HARRIS, J.
HUGHES, E.J. *see* HUSSEY, G.D.
HUSSEY, G.D., WATKINS, M.L.V., GODDARD, E.A., GOTTSCHALK, S., HUGHES, E.J., ILONI, K., KIBEL, M.A. & RESS, S. R. Neonatal mycobacterial specific cytotoxic T-lymphocyte and cytokine profiles in response to district BCG vaccination strategies, 314

IGIETSEME, J.U. *see* MOORE, T.
IKEGAYA, A. *see* IMAI, Y.
ILONI, K. *see* HUSSEY, G.D.
IMAI, Y., FUKUI, T., IKEGAYA, A., ISHIKAWA, T., ONO, Y. & KUROHANE, K. Lack of Shiga-like toxin binding sites in germinal centers of mouse lymphoid tissues, 509
IMANAKA, T. *see* NAGATA, T.
ISHIDA, H. *see* GALUSTIAN, C.
ISHIKAWA, T. *see* IMAI, Y.
IWAKURA, Y. *see* KAKUTA, S.
IZUMIYA, Y. *see* SHIMOJIMA, M.

JENKINSON, E.J. *see* HARE, K.J.
JEON, K.-I. *see* JEONG, J.-Y.
JEONG, J.-Y., CHOI, J.W., JEON, K.-I. & JUE, D.-M. Chloroquine decreases cell-surface expression of tumour necrosis factor receptors in human histiocytic U-937 cells, 83
JUE, D.-M. *see* JEONG, J.-Y.

KAKUTA, S., TAGAWA, Y.-I., SHIBATA, S., NANNO, M. & IWAKURA, Y. Inhibition of B16 melanoma experimental metastasis by interferon-$\gamma$ through direct inhibition of cell proliferation and activation of antitumour host mechanisms, 92
KANG, D. *see* NAGATA, T.
KATO, K. *see* SHIMOJIMA, M.
KATZAV, S. *see* BAR-SHAVIT, R.
KHAN, A.I., HASKARD, D.O., MALHOTRA, R. & LANDIS, R. C. Identification and characterization of L-selectin ligands in porcine lymphoid tissues, 441
KIBEL, M.A. *see* HUSSEY, G.D.
KIMBER, I. *see* CUMBERBATCH, M.
KISHI, H. *see* NAGATA, T.
KISHIHARA, K. *see* UCHIYAMA, H.
KISO, M. *see* GALUSTIAN, C.
KLEIN, G. *see* KUTLEŠA, S.
KOHANAWA, M. *see* DIAO, H.
KOSS, M. *see* HARRIS, J.
KUROHANE, K. *see* IMAI, Y.
KUTLEŠA, S., SILER, U., SPEISER, A., WESSELS, J.T., VIRTANEN, I., ROUSSELLE, P., SOROKIN, L.M., MÜLLER, C.A. & KLEIN, G. Developmentally regulated interactions of human thymocytes with different laminin isoforms, 407

LAAKKO, T. & FRACKER, P. Rapid changes in the lymphopoietic and granulopoietic compartments of the marrow caused by stress levels of corticosterone, 111
LANDIS, R.C. *see* KHAN, A.I.
LEE, T. *see* CAULFIELD, J.
LEE, W.-W., NAM, K.-H., TERAO, K. & YOSHIKAWA, Y. Age-related telomere length dynamics in peripheral blood mononuclear cells of healthy cynomolgus monkeys measured by Flow FISH, 458
LI, J.-H., ROSEN, D., SONDEL, P. & BERKE, G. Immune privilege and FasL: two ways to inactivate effector cytotoxic T lymphocytes by FasL-expressing cells, 267
LIU, Q.L. *see* NAGATA, T.
LLOYD, C. Chemokines in allergic lung inflammation, 144
LOGAN, G.J., SMYTH, C.M.F., EARL, J.W., ZAIKINA, I., ROWE, P.B., SMYTHE, J.A. & ALEXANDER, I.E. HeLa cells cocultured with peripheral blood lymphocytes acquire an immuno-inhibitory phenotype through up-regulation of indoleamine 2,3-dioxygenase activity, 478
LOGAN, M.K. *see* CURNOCK, A.P.
LONDEI, M. *see* OSMAN, M.
LUKEY, P.T. *see* FENHALLS, G.

MA, Y. *see* CHEN, Y.
MACHUGH, N. *see* SATHIYASEELAN, T.
MALHOTRA, R. *see* KHAN, A.I.
MAOZ, M. *see* BAR-SHAVIT, R.
MARCHALONIS, J.J. *see* ROBEY, I.F.
MARTINEZ-A, C. *see* OUTRAM, S.V.
MATHERS, K. *see* SMITH, C.A.
MATSUDA, T. *see* NAGATA, T.
MBAWUIKE, I.M. *see* ZHANG, Y.
MCCARTHY, K. M. *see* VU, Q.
MCCORMACK, J.M. *see* VU, Q.
MCGEE, D.W. *see* MILLER, T.L.
MCGUIRE, T.C. *see* FRASER, D.G.
MERIDA, I. *see* OUTRAM, S.V.
MILLER, H.R.P. & PEMBERTON, A. D. Tissue-specific expression of mast cell granule serine proteinases and their role in inflammation in the lung and gut, 375
MILLER, T.L. & MCGEE, D. W. Epithelial cells respond to proteolytic and non-proteolytic detachment by enhancing interleukin-6 responses, 101
MINAGAWA, A. *see* DIAO, H.
MINAGAWA, R. *see* UCHIYAMA, H.
MIYAZAKI, J.-I. *see* YAMAKAMI, K.
MIYAZAWA, T. *see* SHIMOJIMA, M.
MONAGHAN, P. *see* HARRIS, J.
MOORE, T., ANANABA, G.A., BOLIER, J., BOWERS, S., BELAY, T., EKO, F.O. & IGIETSEME, J.U. Fc receptor regulation of protective immunity against *Chlamydia trachomatis*, 213
MUELLER, C. Tumour necrosis factor in mouse models of chronic intestinal inflammation, 1
MÜLLER, C.A. *see* KUTLEŠA, S.
MULLER, S. *see* BEIGNON, A.-S.
MURAGUCHI, A. *see* NAGATA, T.

NAGATA, T., KISHI, H., LIU, Q.L., MATSUDA, T., IMANAKA, T., TSUKADA, K., KANG, D. & MURAGUCHI, A. The regulation of DNAse activities in subcellular compartments of activated thymocytes, 399
NAIMAN, B. *see* SATHIYASEELAN, S.
NAIR, S. *see* FAIOLA, B.
NAKAJIMA, Y. *see* DIAO, H.
NAKAMURA, K. *see* SHIMOJIMA, M.
NAKANE, A. *see* DIAO, H.
NAM, K.-H. *see* LEE, W.-W.
NANNO, M. *see* KAKUTA, S.
NIELSON, C.P. & WINGETT, D. Endothelial cell and cAMP regulation of T-cell CD40 ligand: relevance of calcium/calmodulin-dependent kinase IV signalling, 430

NISHIMURA, Y. *see* SHIMOJIMA, M.
NITTA, Y. *see* YAMAKAMI, K.
NOMOTO, K. *see* UCHIYAMA, H.
NOURI-ARIA, K.T. *see* WACHHOLZ, P.R.

O'CONNELL, J. Fas ligand and the fate of antitumour cytotoxic T lymphocytes, 263
OAKS, J.L. *see* FRASER, D.G.
ONO, Y. *see* IMAI, Y.
OSMAN, M., TORTORELLA, M., LONDEI, M. & QUARATINO, S. Expression of matrix metalloproteinases and tissue inhibitors of metalloproteinases define the migratory characteristics of human monocyte-derived dendritic cells, 73
OUT, T.A., WANG, S.-Z., RUDOLPH, K. & BICE, D. E. Local T-cell activation after segmental allergen challenge in the lungs of allergic dogs, 499
OUTRAM, S.V., CROMPTON, T., MERIDA, I., VARAS, A. & MARTINEZ-A, C. Diacylgylcerol kinase α activity promotes survival of $CD4^+8^+$ double positive cells during thymocyte development, 391

PACHECO, S. *see* ZHANG, Y.
PARTIDOS, C.D. *see* BEIGNON, A.-S.
PEMBERTON, A.D. *see* MILLER, H.R.P.
PHIPPS, R.P. *see* HARRIS, S.G.
PONGRACZ, J. *see* HARE, K.J.
PRIOR, T.K. *see* HINKLEY, K.S.

QUARATINO, S. *see* OSMAN, M.

RADZUN, H.-J. *see* SORURI, A.
REDECKE, V. *see* HÄCKER, G.
REMMERSWAAL, E.B.M. *see* RENTENAAR, R.J.
RENTENAAR, R.J., VOSTERS, J.L.G., VAN DIEPEN, F.N.J., REMMERSWAAL, E.B.M., VAN LIER, R.A.W. & TEN BERGE, I. J. M. Differentiation of human alloreactive $CD8^+$ T cells *in vitro*, 278
RESS, S.R. *see* HUSSEY, G.D.
RIGGS, J.E. *see* HINKLEY, K.S.
ROBEY, I.F., SCHLUTER, S.F., AKPORIAYE, E., YOCUM, D.E. & MARCHALONIS, J. J. Human monoclonal natural autoantibodies against the T-cell receptor inhibit interleukin-2 production in murine T cells, 419
ROBINSON, J.H. & DELVIG, A. A. Diversity in MHC class II antigen presentation, 252
ROSEN, D. *see* LI, J.-H.
ROTONDO, D. & DAVIDSON, J. Prostaglandin and PPAR control of immune cell function, 20
ROUSSELLE, P. *see* KUTLEŠA, S.
ROWE, P.B. *see* LOGAN, G.J.
RUDOLPH, K. *see* OUT, T.A.

SATHIYASEELAN, T., NAIMAN, B., WELTE, S., MACHUGH, N., BLACK, S.J. & BALDWIN, C. L. Immunological characterization of a γδ T-cell stimulatory ligand on autologous monocytes, 181
SATO, T. *see* DIAO, H.
SCHLUTER, S.F. *see* ROBEY, I.F.
SCHNEEBERGER, E.E. *see* VU, Q.
SCHROEDER, C. *see* SMITH, C.A.
SCHUURMAN, J. *see* AALBERSE, R.C.H.
SCHWEYER, S. *see* SORURI, A.
SHIBATA, S. *see* KAKUTA, S.
SHIMOJIMA, M., NISHIMURA, Y., MIYAZAWA, T., KATO, K., NAKAMURA, K., IZUMIYA, Y., AKASHI, H. & TOHYA, Y. A feline CD2 homologue interacts with human red blood cells, 360
SILER, U. *see* KUTLEŠA, S.
SKINNER, A. *see* SMITH, C.A.
SMITH, C.A., GRAHAM, C.M., MATHERS, K., SKINNER, A., HAY, A.J., SCHROEDER, C. & THOMAS, D. B. Conditional ablation of T-cell development by a novel viral ion channel transgene, 306
SMYTH, C.M.F. *see* LOGAN, G.J.
SMYTHE, J.A. *see* LOGAN, G.J.
SNETKOV, V. *see* CAULFIELD, J.
SONDEL, P. *see* LI, J.-H.
SOROKIN, L.M. *see* KUTLEŠA, S.
SORURI, A., SCHWEYER, S., RADZUN, H.-J. & FAYYAZI, A.. Mycobacterial antigens induce apoptosis in human purified protein derivative-specific αβ T lymphocytes in a concentration-dependent manner, 222
SPEISER, A. *see* KUTLEŠA, S.
STEVENS, L. *see* FENHALLS, G.
STOLL, M. *see* GALUSTIAN, C.
SUGIMACHI, K. *see* UCHIYAMA, H.
SWAIN, M.G. *see* AJUEBOR, M.N.
SWITZER, K.C. *see* ZHANG, Y.

TADAKUMA, T. *see* YAMAKAMI, K.
TAGAWA, Y.-I. *see* KAKUTA, S.
TAYLOR, G. *see* HARRIS, J.
TEN BERGE, I.J.M. *see* RENTENAAR, R.J.
TERAO, K. *see* LEE, W.-W.
THOMAS, D.B. *see* SMITH, C.A.
TILL, S.J. *see* WACHHOLZ, P.R.
TOHYA, Y. *see* SHIMOJIMA, M.
TORTORELLA, M. *see* OSMAN, M.
TSUKADA, K. *see* NAGATA, T.

UCHIYAMA, H., KISHIHARA, K., MINAGAWA, R., HASHIMOTO, K., SUGIMACHI, K. & NOMOTO, K. Crucial Fas–Fas ligand interaction in spontaneous acceptance of hepatic allografts in mice, 450

VAN DIEPEN, F.N.J. *see* RENTENAAR, R.J.
VAN LIER, R.A.W. *see* RENTENAAR, R.J.
VARAS, A. *see* OUTRAM, S.V.
VERHOEF, A. *see* WACHHOLZ, P.R.
VIRTANEN, I. *see* KUTLEŠA, S.
VOSTERS, J.L.G. *see* RENTENAAR, R.J.
VU, Q., MCCARTHY, K.M., MCCORMACK, J.M. & SCHNEEBERGER, E. E. Lung dendritic cells are primed by inhaled particulate antigens, and retain MHC class II/antigenic peptide complexes in hilar lymph nodes for a prolonged period of time, 488

WACHHOLZ, P.R., NOURI-ARIA, K.T., WILSON, D.R., WALKER, S.M., VERHOEF, A., TILL, S.J. & DURHAM, S. R. Grass pollen immunotherapy for hayfever is associated with increases in local nasal but not peripheral Th1. Th2 cytokine ratios, 56
WALKER, S.M. *see* WACHHOLZ, P.R.
WANG, S.-Z. *see* OUT, T.A.
WANG, S. *see* YANG, X.
WANG, Y. *see* ZHANG, Y.
WARD, S.G. *see* CURNOCK, A.P.
WATKINS, M.L.V. *see* HUSSEY, G.D.
WATSON, M. L. Chemokines – linking receptors to response, 121
WELTE, S. *see* SATHIYASEELAN, T.
WERLING, D. *see* HARRIS, J.
WESSELS, J.T. *see* KUTLEŠA, S.
WILSON, D.R. *see* WACHHOLZ, P.R.
WINGETT, D. *see* NIELSON, C.P.

YAMAKAMI, K., AKAO, S., TADAKUMA, T., NITTA, Y., MIYAZAKI, J.-I. & YOSHIZAWA, N. Administration of plasmids expressing interleukin-4 and interleukin-10 causes BALB/c mice to induce a T helper 2-type response despite the expected T helper 1-type response with a low-dose infection of *Leishmania major*, 515

YANG, J. *see* YANG, X.
YANG, X., FAN, Y., WANG, S., HAN, X., YANG, J., BILENKI, L. & CHEN, L. Mycobacterial infection inhibits established allergic inflammatory responses via alteration of cytokine production and vascular cell adhesion molecule-1 expression, 336
YIMIN, H. *see* DIAO, H.
YOCUM, D.E. *see* ROBEY, I.F.
YONGJUN, Y. *see* BAR-SHAVIT, R.
YOSHIKAWA, Y. *see* LEE, W.-W.
YOSHIZAWA, N. *see* YAMAKAMI, K.

ZAIKINA, I. *see* LOGAN, G.J.
ZHANG, Y., PACHECO, S., ACUNA, C.L., SWITZER, K.C., WANG, Y., GILMORE, X., HARRIMAN, G.R. & MBAWUIKE, I. M. Immunoglobulin A-deficient mice exhibit altered T helper 1-type immune responses but retain mucosal immunity to influenza virus, 286

# Subject Index


Activation-induced cell death (AICD):
  by mycobacterial antigens, 222
  PPD-induced, 222
  T cells, 222
Allergic inflammation:
  inhibited by VCAM-1, 337
Antigen presentation:
  CTL induction by DCs, 47
Antigen-presenting cells:
  bovine, 190
  calveolin expression, 190
Autoantibodies:
  against TCR, 419
  human, 419
  natural, 419

B-1b B cells:
  age-dependent increase, 196
  in SCID mice, 196
  peritoneal, 196
BCG vaccine:
  cellular immunity, 314
  neonates, 314
  Th1 cytokines, 314

Calveolin expression:
  bovine APC, 190
Caspase:
  caspase-activated deoxyribonuclease (CAD), 399
  role in thymocyte apoptosis, 399
CD2:
  cDNA, 361
  feline, 361
  human, 361
$CD4^+CD8^+$ double-positive cells:
  lipid mediators, 391
  promotion of survival, 391
  role of diacylglycerol kinase in survival, 391
CD8 cells:
  alloreactive, 278
  differentiation, 278
  human, 278
CD40 L:
  regulation, 430
  role in endothelial cell regulation, 430
Chemokine receptor expression:
  effect of dexamethasone, 155
  human, 155
  monocytes, 155
  regulated by $CD3^+$ T-cell factors, 155
Chemokines:
  airway inflammation, 144
  allergy, 144
  gastrointestinal disease, 137
  inflammatory bowel disease, 137
  ligands, 121
  liver disease, 137
  P13 K activation, 125
  receptors, 121, 137
  signalling, 125
  Th2 cells, 144
*Chlamydia trachomatis*:
  protective immunity, 213
  Th1 response, 213
  vaccine efficacy, 213

Costimulation:
  synergy between endothelial cells and cAMP, 430
CpG immunization:
  bare skin, 204
  influenza virus, 204
CpG-DNA:
  activation of immune system, 245
  bacterial, 245
CTLA-4:
  role in oral tolerance, 171
Cytokine:
  production during *M. tuberculosis* infection, 326

Dendritic cells:
  antigen presentation capacity in lung, 488
  class II epitopes, 47
  cytotoxic T-cell priming, 47
  hilar lymph nodes, 488
  human, 73
  MHC class II expression, 488
  lung, 488
  role of MMP and TIMP in migration, 73

Endothelial cells:
  CD40 L expression, 430
  costimulation, 430
Epithelial cells:
  cell detachment, 101
  IL-6, 101

Fas–Fas ligand interaction:
  in liver transplantation, 450
  murine allograft, 450
Fas L:
  CTL apoptosis, 263
  murine, 263
  role in antitumour response, 263
Fas L expression:
  in CTLs, 267
  role in immune privilege, 267
  tumours, 267

GvH disease:
  eosinophil, 63
  role of IFN $\gamma$ in pathogenesis, 63
$\gamma\delta$ T cells:
  bovine, 181
  MLR, 181
  stimulation by monocytes, 181

Haemopoeisis:
  role of corticosterone, 111
  stress, 111
Hayfever:
  immunotherapy, 56
HeLa cells:
  IDO activity, 478
  immunoinhibitory phenotype, 478

IBD:
  murine, 1
  role of TNF, 1
IgE-mediated allergy:
  immune complex, 9
  role of IgG4, 9

IgG4:
bispecificity, 9
chimeric antibody, 9
IL-1β:
epidermal, 466
role in LC migration, 466
IL-6:
production by intestinal epithelial cells, 101
IL-10:
B16-melanoma proliferation, 231
mice–melanoma model, 231
promotes melanoma growth, 231
Immune privilege:
AICD, 267
Fas L-expressing tumours, 267
Immunomodulation:
CpG immunization, 204
influenza priming, 204
of virus-specific $CD4^+$ T cells, 204
Immunotherapy:
hayfever, 56
nasal cytokine, 56
Th1 : Th2 cytokine ratios, 56
Inflammation:
chemokine receptor role, 137
chemokine role, 137
Influenza:
virus-specific $CD4^+$ T cells, 204
Influenza virus:
IgA deficiency, 286
immunization, 286
mucosal immunity, 286
Innate immunity:
bacterial CpG-DNA, 245

Langerhan cell migration:
ageing, 466
allergen-induced, 466
murine, 466
role of IL-1β, 466
*Leishmania major* infection:
role of IL-4 and IL-10, 515
Th2 response, 515
Lethal transgene:
T cells, 306
thymopoiesis, 306
Lipid mediators:
$CD4^+CD8^+$ cells, 391
L-selectin:
porcine, 441
characterization, 441

*M. tuberculosis*:
CD8 cells, 326
cytokines, 326
human, 326
Mast cell:
proteinases, 375
Melanoma:
antitumour mechanisms, 92
B16 melanoma, 92
cell-mediated immunity, 92
IL-10-induced proliferation, 231
murine, 231
role of IFN γ, 92
Metalloproteinases:
human, 73
role in DC migration, 73
MHC class II presentation:
diverse peptide repertoire, 252
Migration:
human DCs, 73
MMP, 73
TIMP, 73
Mycobacterial:
neonatal immunity, 314
specific CTL, 314
Mycobacterial infection:
inhibits allergic inflammation, 337
role of VCAM-1, 337

Notch-1 signalling:
$CD4^+$ $CD8^+$ thymocytes, 163

Oral tolerance:
induction, 171
OVA, 171
role of CTLA-4, 171

PAMPs (pathogen-associated molecular patterns):
role in innate immunity, 245
stimulatory potential, 245
PBL:
coculture with HeLa cells, 478
IDO activity, 478
human, 478
Peroxisome proliferator-activated receptors (PPARs):
immunoregulatory role, 20
Prostaglandins:
1, 20
control of the immune system, 20
role in apoptosis, 23
Proteinases:
chymase, 375
mast cell granule, 375
role in gut/lung inflammation, 375
tryptase, 375

*S. pyogenes*:
murine infection, 345
role of TNF α, 345
SCID mice:
aging, 196
B-lymphocyte generation, 196
Selectins:
interactions, 351
interactions with lipid-linked sulpho-tyrosine probe, 351
therapeutic inhibitor designs, 351
Shiga toxin:
binding in germinal centres, 509
murine CD77, 509

T lymphocytes:
apoptosis, 23
prostaglandin receptor expression, 23
T-cell activation:
allergen challenge, 499
allergic asthma, 499
dog, 499
lung, 499
T-cell signalling pathways:
protease-activated receptor (PAR), 35
role of integrin activation, 35
Var 1, 35
TCR:
autoantibodies, 419
human, 419
protein tyrosine kinase, 369
signalling, 369
Telomere length:
in T-cell subsets, 458
measurement, 458
monkey PBMC, 458
Th1:
equine, 295
epitopes in equine lentivirus, 295

Th1 : Th2 cytokines:
hayfever immunotherapy, 56
nasal, 56
Th2 cytokines:
persistence in *L. major* infection, 515
Thymocyte:
apoptosis, 399
caspases, 399
human, 407
laminin expression, 407
negative selection, 163
positive selection, 163
resistance to apoptosis, 163
Thymocyte negative selection:
independent of Notch-1 signalling, 163
Thymopoiesis:
ion channel activity, 306
lineage-specific cell ablation, 306
TNF receptor expression:
chloroquine suppresses, 83
human, 83
U-937 cells, 83
TNF:
intestinal inflammation, 1
murine IBD, 1
role in colitis, 1
TNF $\alpha$:
role in streptococcal disease, 345

VCAM-1:
role in allergic inflammation, 337